

HIV/AIDS CASES
ALABAMA
Reported 1982 through November 12, 2004

| Exposure Categories | AIDS | % | HIV | % |
|----------------------------|-------------|----------|-------------|----------|
| MSWM | 3595 | 46.1% | 2119 | 33.8% |
| IDU | 1104 | 14.2% | 600 | 9.6% |
| MSWM/IDU | 621 | 8.0% | 254 | 4.0% |
| Transfusion | 89 | 1.1% | 50 | 0.8% |
| Hemophiliac | 68 | 0.9% | 27 | 0.4% |
| Heterosexual Contact | 1246 | 16.0% | 1416 | 22.6% |
| Perinatal | 72 | 0.9% | 34 | 0.5% |
| Other/Not Reported | 1002 | 12.9% | 1773 | 28.3% |
| Race/Gender | | | | |
| Af-Am Female | 1162 | 14.9% | 1415 | 22.6% |
| Af-Am Male | 3605 | 46.2% | 2569 | 41.0% |
| White Female | 343 | 4.4% | 414 | 6.6% |
| White Male | 2586 | 33.2% | 1674 | 26.7% |
| Other*/Unknown Female | 19 | 0.2% | 49 | 0.8% |
| Other*/Unknown Male | 82 | 1.1% | 152 | 2.4% |
| Mortality Status | | | | |
| Living | 4132 | 53.0% | 5948 | 94.8% |
| Deceased | 3665 | 47.0% | 325 | 5.2% |
| Agegroup | | | | |
| 0 to 4 | 64 | 0.8% | 28 | 0.4% |
| 5 to 12 | 12 | 0.2% | 13 | 0.2% |
| 13 to 19 | 70 | 0.9% | 318 | 5.1% |
| 20 to 24 | 410 | 5.3% | 1002 | 16.0% |
| 25 to 29 | 1186 | 15.2% | 1201 | 19.1% |
| 30 to 34 | 1695 | 21.7% | 1235 | 19.7% |
| 35 to 39 | 1592 | 20.4% | 976 | 15.6% |
| 40 to 44 | 1142 | 14.6% | 655 | 10.4% |
| 45 to 49 | 738 | 9.5% | 420 | 6.7% |
| 50 and over | 888 | 11.4% | 398 | 6.3% |
| Unknown | 0 | 0.0% | 27 | 0.4% |
| TOTAL | 7797 | | 6273 | |

* Due to low morbidity in these groups and space constraints in this report, Hispanic, Asian/Pacific and Native American races are included as "Other" races.

HIV count does not include perinatally exposed infants who have either seroreverted or whose infection status has yet to be determined.