



Anal fissure

Faeces is temporarily stored in the rectum and expelled from the body via the anus. An anal fissure is a tear or split in the lining of the anus (anal mucosa). The symptoms include pain when passing a bowel motion and bright red blood from the anus. The problem is common in children younger than one year, and affects around eight out of 10 babies. A person's susceptibility to anal fissures tends to decline with age. Common causes in adults include constipation and trauma to the anus (such as a difficult childbirth). Around half of all cases heal by themselves, with proper self-care and avoidance of constipation. However, healing can be a problem if the pressure of passing bowel motions constantly reopens the fissure. Treatment options include surgery.

Symptoms

The symptoms of an anal fissure include:

- Anal pain
- Pain when passing a motion and for some time afterwards
- Bright red blood from the anus
- Blood on the surface of stools
- Blood smears on the toilet paper.

The anus explained

The anus permits bowel continence, which is voluntary control over the passage of bowel motions. Waste is massaged down the length of the large intestine by waves of muscular contractions (peristalsis). Once the excess water is removed, the waste is temporarily stored in the rectum. The rectum joins to the anus, which is a short tube containing a muscular ring (sphincter) that can be opened at will to allow the expulsion of faeces. The membranous lining of the anus is called the anal mucosa. An anal fissure is a tear in the anal mucosa.

A range of causes

Some of the causes of an anal fissure include:

- Chronic constipation
- Passing a dry, hard stool
- Rough or excessive wiping of the anus after passing a motion
- Diarrhoea
- Inflammation of the anus and rectum
- Crohn's disease
- Scratching (as a reaction to pinworm infection, for example)
- Anal injury
- Pregnancy
- Childbirth.

Possible complications

Anal fissures aren't associated with more serious diseases, such as bowel cancer. Some of the possible complications of an anal fissure can include:

- **Chronic anal fissure** – the tear fails to heal. Over time, this can cause extensive scar tissue at the site of the fissure (sentinel pile).
- **Anal fistulas** – abnormal 'tunnels' join the anal canal to the skin around the anus and sometimes to other parts of the bowel.
- **Anal stenosis** – the anal canal becomes abnormally narrowed either due to spasm of the anal sphincter or contraction of the resultant scar tissue.

Diagnosis methods

An anal fissure is diagnosed using a number of tests including:

- Physical examination
- Inspection of the anus and rectum with a slender instrument (anoscope).

Treatment options

Medical treatment for an anal fissure may include:

- Pain-killing medications
- Laxatives
- Anaesthetic creams
- Nitroglycerin creams
- Surgery.

Surgical correction

Severe anal fissures need to be surgically corrected. The fissure and associated scar tissue is removed. Sometimes, a thin slice of muscle from the anal sphincter is also removed, as this helps the wound to better heal. (Cutting and suturing this muscle doesn't interfere with the patient's sphincter control.) The operation, called a lateral internal sphincterotomy, can be performed under local anaesthetic. Around nine out of 10 patients will never experience another anal fissure.

Self-help treatment options

Be guided by your health care professional, but general suggestions include:

- Apply petroleum jelly to the anus.
- See your chemist for advice on ointments specific for anal pain.
- Take regular sitz baths, which involve sitting in a shallow bath of warm water for around 20 minutes.
- Use baby wipes instead of toilet paper.
- Shower or bathe after every bowel motion.
- Drink six to eight glasses of water every day.

See over ...



Prevention suggestions

Many cases of anal fissure are caused by chronic constipation. Suggestions include:

- Eat a high fibre diet
- Drink plenty of water to help soften stools
- Consider using a fibre supplement (such as Metamucil)
- Make sure to wipe gently after going to the toilet.

Where to get help

- Your doctor
- Chemist

Things to remember

- Faeces is temporarily stored in the rectum and expelled from the body via the anus.
- An anal fissure is a tear or split in the lining of the anus (anal mucosa).
- The symptoms include pain and bright red blood from the anus.
- Treatment options include laxatives and surgery.
- Self-help suggestions include switching to a high fibre diet and drinking plenty of water.

This page has been produced in consultation with, and approved by, the North East Valley Division of General Practice. The Better Health Channel is part of the Department of Human Services, Victoria