



Effect of breast feeding on intelligence in children: prospective study, sibling pairs analysis, and meta-analysis

Geoff Der, G David Batty and Ian J Deary

BMJ 2006;333:945-; originally published online 4 Oct 2006;
doi:10.1136/bmj.38978.699583.55

Updated information and services can be found at:
<http://bmj.com/cgi/content/full/333/7575/945>

These include:

Data supplement

"Further details"

<http://bmj.com/cgi/content/full/bmj.38978.699583.55/DC1>

References

This article cites 29 articles, 8 of which can be accessed free at:
<http://bmj.com/cgi/content/full/333/7575/945#BIBL>

2 online articles that cite this article can be accessed at:
<http://bmj.com/cgi/content/full/333/7575/945#otherarticles>

Rapid responses

14 rapid responses have been posted to this article, which you can access for free at:

<http://bmj.com/cgi/content/full/333/7575/945#responses>

You can respond to this article at:

<http://bmj.com/cgi/eletter-submit/333/7575/945>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top left of the article

Topic collections

Articles on similar topics can be found in the following collections

[Other Psychiatry](#) (904 articles)

[Other nutrition and metabolism](#) (1272 articles)

[Breastfeeding and infant nutrition](#) (160 articles)

[Children](#) (1805 articles)

Notes

To order reprints follow the "Request Permissions" link in the navigation box

To subscribe to *BMJ* go to:

<http://resources.bmj.com/bmj/subscribers>

Effect of breast feeding on intelligence in children: prospective study, sibling pairs analysis, and meta-analysis

Geoff Der, G David Batty, Ian J Deary

Abstract

Objective To assess the importance of maternal intelligence, and the effect of controlling for it and other important confounders, in the link between breast feeding and children's intelligence.

Design Examination of the effect of breast feeding on cognitive ability and the impact of a range of potential confounders, in particular maternal IQ, within a national database. Additional analyses compared pairs of siblings from the sample who were and were not breast fed. The results are considered in the context of other studies that have also controlled for parental intelligence via meta-analysis.

Setting 1979 US national longitudinal survey of youth.

Subjects Data on 5475 children, the offspring of 3161 mothers in the longitudinal survey.

Main outcome measure IQ in children measured by Peabody individual achievement test.

Results The mother's IQ was more highly predictive of breastfeeding status than were her race, education, age, poverty status, smoking, the home environment, or the child's birth weight or birth order. One standard deviation advantage in maternal IQ more than doubled the odds of breast feeding. Before adjustment, breast feeding was associated with an increase of around 4 points in mental ability. Adjustment for maternal intelligence accounted for most of this effect. When fully adjusted for a range of relevant confounders, the effect was small (0.52) and non-significant (95% confidence interval -0.19 to 1.23). The results of the sibling comparisons and meta-analysis corroborated these findings.

Conclusions Breast feeding has little or no effect on intelligence in children. While breast feeding has many advantages for the child and mother, enhancement of the child's intelligence is unlikely to be among them.

Introduction

Since 1929 many studies have examined the link between breast feeding and intelligence, most finding higher IQ scores among children who were breast fed.¹ The few randomised controlled trials were confined to preterm infants,²⁻⁴ whereas studies of individuals born at term were observational in design. Inference from the observational studies is hampered by confounding: there are fundamental differences between mothers who choose to breast feed and those who do not. Studies that do not control adequately for confounders may mistake residual confounding for a real effect of breast feeding.

Many potential confounding variables have been identified, including duration of breast feeding, sex, maternal history of

smoking, maternal age, maternal intelligence, maternal education, maternal training, paternal education, race or ethnicity, socioeconomic status, family size, birth order, birth weight, gestational age, and childhood experiences.⁵ Those singled out as particularly important include socioeconomic status, maternal education, and birth weight⁶ or socioeconomic status/parental education and stimulation of the child.⁷ In contrast, maternal intelligence is relatively overlooked as a potential confounder. This is surprising given the heritability of intelligence⁸ and the known association of maternal intelligence with both the initiation and duration of breast feeding.⁹

We examined the relation between breast feeding and intelligence and assessed the role of maternal IQ and other covariates in generating the association. We took both a conventional approach to control for confounders and an alternative approach using sibling comparison analysis. Used recently to assess the benefits of breast feeding,¹⁰ this approach has the considerable advantage of controlling for many confounding factors without having to measure them. Any factor that is the same for both members of a pair of siblings is automatically and fully controlled for. Thus, the method implicitly controls for parental intelligence.

Methods

We used data from the US national longitudinal survey of youth 1979 (NLSY79).¹¹ This is a population based sample of 12 686 young people (6283 female) aged 14 to 22 when first interviewed in 1979 who were then interviewed annually until 1994 and biennially thereafter. Since 1986 the children of the women in the survey have also been assessed biennially. The database for the children is referred to as the NLSY79 child and young adult sample ("young adult" because some of the children are now of adult age).¹² As preterm babies are known to have different nutritional needs¹³ we excluded children born before the 35th week and those who weighed less than 2500 g at birth. We also excluded those born before 1979, when the survey began.

Measures

Children's cognitive ability—The Peabody individual achievement test (PIAT) was administered to children between the ages of 5 and 14 biennially from 1986 to 2002. Children were tested repeatedly if they fell within the age range in test years. We used the PIAT total scores, as well as the individual component scores for mathematics, reading comprehension, and reading recognition. To aid comparison of effect sizes across measures and with the results of other studies, we standardised all outcomes to a mean of 100 and standard deviation of 15.

Research

Breast feeding—Women who had had a child since the previous interview were asked whether they breast fed the child at all and, if so, how old the child was when they stopped breast feeding. In most cases this information would have been obtained within a year of the birth. Even so, reports on duration of breast feeding were probably less reliable than reports of whether a child was breast fed or not.

Control variables—We chose control variables on the basis of associations with breast feeding or childhood cognitive development in the survey,^{14 15} other studies of breast feeding and cognitive development,^{9 16 17} and the most recent reviews of the subject.^{5-7 18}

Child's environment—The short form of the home observation for measurement of the environment scale (HOME-SF)¹⁹ was completed at each assessment, and we used the cognitive stimulation and emotional support subscales.

Child demographics—Sex, age, gestation (weeks), birth weight, and birth order were also used in the models.

Maternal characteristics—Maternal cognitive ability was measured with the armed forces qualification test (AFQT).¹¹ This is derived from the arithmetic reasoning, word knowledge, paragraph comprehension, and mathematical knowledge sections of the Armed Services vocational aptitude battery. It was administered in 1980. Level of education was classified as the highest grade completed by the mother at the time of the child's birth. Race was classified as Hispanic, black, or non-Hispanic/non-black. The latter includes white people and others who did not identify themselves as black or Hispanic. Poverty status indicates whether the household income was below the poverty threshold at the birth of the child. We also included the mother's age at the birth of the child plus a binary variable to indicate whether the mother smoked during the pregnancy.

Statistical analysis

Linkage of the data from the survey with the child and young adult sample resulted in a dataset that contained one or more assessments for each child and one or more children per mother. This can be viewed as a hierarchical structure of assessments nested within children and children nested within mothers. The multiple assessments made on each child are unlikely to be statistically independent, nor are the data for siblings. To accommodate this lack of independence we used random effects models, which have the advantage of being able to use all available assessments for each child and all children, whether they are siblings or not. The statistical background for this use of random effects models is given elsewhere,^{20 21} and tutorial articles showing analyses similar to those conducted here are also available.²²⁻²⁵

All analyses were carried out separately for PIAT total score and the three subscales: mathematics, reading recognition, and reading comprehension. Initial analysis assessed the association between each of the potential confounders and breast feeding. The main analyses estimated the effect of breast feeding on cognitive ability; examined the extent to which each of the confounders, individually, attenuated the effect; and then examined the attenuation of multiple confounders.

To use the random effects models we structured data so that each occasion when measurements were done was a separate record in the data file. Each child then has one or more records depending on the number of PIAT assessments completed. Data that were not assessed repeatedly, or do not change with time, were attached to this record, including, for example, information about the mother, the pregnancy, and the child's birth. Each record was also indexed with some time or occasion variable. We

used the child's age at the time of the assessment, centred at 10 years. To control for the correlation between repeated measures and between siblings, the models included random intercepts, random slopes in age (the child's), and intercept-slope covariance, both at the child level and at the level of the mother. Thus there were six random effects in each of the models. All models also included a fixed effect of child's age. The models were estimated by restricted maximum likelihood with proc mixed in SAS version 9.1.

As the sample contained a large number of siblings, it was also well suited to examining the same question with sibling comparison analysis. A simple test of the null hypothesis of no effect of breast feeding can be made by identifying pairs of siblings in which one sibling was breast fed and the other was not, subtracting the PIAT scores of the sibling who was not breast fed from those of the one who was, and testing whether the resulting differences have a mean of zero. The same can be done for sibling pairs that are discordant for duration of breast feeding, subtracting the scores of the sibling with the shorter duration of breast feeding.

We considered the results of both types of analysis in the context of other comparable studies via meta-analysis (comprehensive meta-analysis program version 2.2.020, www.meta-analysis.com). There are no agreed standards of study quality in this area, and the four recent reviews identified all used different criteria, resulting in sets of studies with relatively little overlap.^{5-7 18} We included only studies that quantified the effect of breastfeeding status on cognitive ability after controlling for parental intelligence among full term infants. We searched Medline (1966 to January 2006), ISI Web of Knowledge, and the British Library's electronic table of contents for any combination of "breast feeding/breast milk/infant feeding" and "cognitive development/intelligence/intellectual development" and supplemented the results with additional studies identified from the reference lists of the four reviews mentioned and from those of individual studies.

Results

Table 1 shows descriptive statistics for the potential confounders, broken down by breastfeeding status. The association of each of them with breastfeeding status is shown as an odds ratio. Children who were breast fed had mothers with higher IQ and with more education and who were older, less likely to be in poverty or to smoke, and more likely to provide a more stimulating and supportive home environment. Hispanic mothers were less likely to breast feed their children and black mothers much less likely. The children who were breast fed were likely to be heavier at birth and earlier in birth order, although this could be a reflection of family size. There was no difference in gestation or the proportion of male infants who were breast fed.

A one standard deviation advantage in mother's IQ more than doubled of the odds of breast feeding. Mother's education had a similar but slightly weaker effect. As gestation and sex were not significantly associated with breast feeding, they were not included in the remaining analyses. For the analyses of PIAT scores we had full data on 16 744 assessments of 5475 children born to 3161 mothers.

Table 2 shows the effect of breast feeding on the cognitive outcomes, both unadjusted and adjusted individually for each of the confounders. Each row gives the effect of breast feeding on the outcome as a regression coefficient (B) with its standard error and P value, adjusted for the confounder listed. The attenuation of the effect size is shown as a percentage of the

Table 1 Association of potential confounders with breast feeding

Confounder	Not breast fed		Breast fed		Odds ratio* (95% confidence interval)	P value
	Mean (SD) or %	No	Mean (SD) or %	No		
Mother's AFQT score	26.24 (22.33)	3001	46.44 (27.76)	2701	2.30 (2.16 to 2.44)	<0.001
Mother's education	11.50 (2.14)	3119	12.79 (2.48)	2819	1.83 (1.73 to 1.95)	<0.001
Mother's age	23.83 (4.50)	3125	25.79 (4.64)	2823	1.55 (1.47 to 1.64)	<0.001
HOME cognitive stimulation	94.09 (16.62)	2844	101.00 (14.26)	2620	1.59 (1.50 to 1.69)	<0.001
HOME emotional support	95.18 (16.54)	2791	100.67 (14.45)	2601	1.44 (1.36 to 1.52)	<0.001
Gestation (weeks)	39.05 (1.58)	3125	39.01 (1.55)	2823	0.97 (0.93 to 1.03)	0.315
Birth weight (g)	3390 (47)	3125	3470 (47)	2823	1.21 (1.15 to 1.27)	<0.001
Birth order	1.97 (1.07)	3125	1.85 (1.02)	2823	0.90 (0.85 to 0.94)	<0.001
Family in poverty†	36.4%	3108	15.8%	2817	0.33 (0.29 to 0.37)	<0.001
Mother smoked in pregnancy†	35.7%	3120	25.6%	2816	0.62 (0.55 to 0.69)	<0.001
Male infant†	51.7%	3125	50.6%	2823	0.96 (0.87 to 1.06)	0.413
Hispanic‡	49.4%	615	50.6%	630	0.72 (0.63 to 0.82)	<0.001
Black‡	74.9%	1264	25.1%	423	0.24 (0.21 to 0.27)	<0.001
Not black/Hispanic‡	41.3%	1246	58.7%	1770	1.00	

AFQT=armed forces qualification test; HOME=home observation of the environment.

*Odds ratios for breast feeding are per 1 SD of mother's AFQT score, education, age, HOME scores, gestation, and birth weight.

†Reference groups are family not in poverty, mother did not smoke in pregnancy, female.

‡Row percentage.

unadjusted figure. The unadjusted effects of breast feeding correspond to an advantage for those breast fed of 4.1 to 4.7. These are comparable with effects reported in other studies.⁵ Adjustment for mother's IQ reduced this advantage by 71% to 75%, and adjustment for mother's education by 34% to 42%. Family poverty, maternal race, maternal age, and HOME stimulation score were important confounders.

Table 3 shows the results of including all the confounders simultaneously. In this case, each row of the table gives the effect of the confounder on the outcome, adjusted for all the others and for the child's age at assessment. The fully adjusted effect of breast feeding averages slightly less than half a point with a range of 0.36 to 0.52. These are small effects, and none is significant even with the large numbers in these analyses. Maternal IQ has the largest independent effect, with maternal education, age, family poverty, HOME stimulation score, and birth order all making independent contributions for most outcomes.

To simulate a study without adjustment for maternal intelligence, but which was otherwise well controlled, we re-ran the models in table 3 omitting maternal IQ. The effects of breast feeding were then at least double those shown in table 3 and all were significant (table 4). Although these effects were still small,

they do show that omitting maternal intelligence can seriously overestimate the effect of breast feeding.

Because observational studies use dose-response relation to infer causality²⁶ we repeated the analysis including data only on those who were breast fed and introducing duration of breast feeding in months (defined as four weeks) to the model (table 4). Although the effects for reading comprehension and PIAT total score were significant, all effects were small considering that the median duration of breast feeding is three months and the 95th centile is 14 months. We repeated the analysis dividing those breast fed into quarters of duration of breast feeding and contrasted each of these with those not breast fed (table 4). The results broadly concur with the previous analysis. There were significant differences for the PIAT total score and the maths and reading comprehension subscales, but only for those in the top quarter of duration of breast feeding. There were also some signs of a departure from linearity in the second quarter.

Sibling pairs analysis

There were 332 sibling pairs discordant for breastfeeding status and 545 discordant for duration of breast feeding. Table 5 shows the mean differences between both groups of siblings for PIAT

Table 2 Effect of breast feeding on cognitive outcomes, unadjusted and adjusted singly for each confounder, in 3161 mothers, 5475 children, and 16 744 assessments. All significant at P<0.001 except where marked

	PIAT-total		Maths*		Reading*		Comprehension*	
	B (SE)	%†	B (SE)	%†	B (SE)	%†	B (SE)	%†
Unadjusted	4.69 (0.38)		4.65 (0.36)		4.09 (0.38)		4.22 (0.36)	
Adjusted for:								
Mother's AFQT score	1.30 (0.36)	72	1.30 (0.34)	72	1.02 (0.37)‡	75	1.21 (0.35)	71
Mother's education	2.95 (0.37)	37	3.06 (0.35)	34	2.38 (0.37)	42	2.69 (0.35)	36
Family in poverty	3.94 (0.38)	16	3.96 (0.36)	15	3.30 (0.37)	19	3.46 (0.35)	18
Mother's age	4.29 (0.38)	9	4.18 (0.36)	10	3.64 (0.38)	11	3.96 (0.36)	6
Mother smoked in pregnancy	4.60 (0.38)	2	4.60 (0.36)	1	3.98 (0.38)	3	4.14 (0.36)	2
HOME cognitive stimulation	4.29 (0.37)	8	4.20 (0.35)	10	3.66 (0.37)	10	3.64 (0.35)	14
HOME emotional support	4.57 (0.38)	3	4.47 (0.36)	4	3.96 (0.38)	3	4.06 (0.36)	4
Birth weight	4.60 (0.38)	2	4.52 (0.36)	3	4.02 (0.38)	2	4.15 (0.36)	2
Birth order	4.55 (0.38)	3	4.57 (0.36)	2	3.94 (0.37)	4	4.01 (0.35)	5
Race	3.65 (0.38)	22	3.30 (0.36)	29	3.32 (0.39)	19	3.34 (0.37)	21

PIAT=Peabody individual achievement test; AFQT=armed forces qualification test; HOME=home observation for measurement of the environment.

*Individual components of PIAT.

†Percentage of unadjusted figure.

‡P=0.006.

Research

Table 3 Mutually adjusted effects of breast feeding and confounders on cognitive outcomes in 3161 mothers, 5475 children, and 16 744 assessments

Confounder	PIAT-total		Maths*		Reading*		Comprehension*	
	B (SE)	P	B (SE)	P	B (SE)	P	B (SE)	P
Breast feeding	0.52 (0.36)	0.149	0.52 (0.34)	0.130	0.36 (0.37)	0.332	0.52 (0.35)	0.134
Mother's AFQT score	4.43 (0.26)	<0.001	3.87 (0.25)	<0.001	3.77 (0.27)	<0.001	3.97 (0.25)	<0.001
Mother's education	1.03 (0.24)	<0.001	1.10 (0.23)	<0.001	0.96 (0.25)	<0.001	0.62 (0.23)	0.007
Family in poverty	-1.72 (0.41)	<0.001	-0.98 (0.39)	0.012	-1.70 (0.42)	<0.001	-1.82 (0.39)	<0.001
Mother's age	0.98 (0.20)	<0.001	0.72 (0.19)	<0.001	1.05 (0.20)	<0.001	0.69 (0.19)	<0.001
Mother smoked in pregnancy	0.08 (0.38)	0.839	0.37 (0.36)	0.305	-0.11 (0.39)	0.771	0.14 (0.36)	0.694
HOME cognitive stimulation	0.83 (0.10)	<0.001	0.78 (0.12)	<0.001	0.79 (0.11)	<0.001	1.13 (0.12)	<0.001
HOME emotional support	0.17 (0.09)	0.072	0.25 (0.11)	0.020	0.15 (0.10)	0.120	0.14 (0.11)	0.200
Birth weight	0.32 (0.16)	0.047	0.40 (0.15)	0.010	0.20 (0.17)	0.234	0.25 (0.16)	0.113
Birth order	-1.54 (0.18)	<0.001	-0.79 (0.17)	<0.001	-1.47 (0.19)	<0.001	-1.69 (0.18)	<0.001
Hispanic	-0.36 (0.52)	0.494	-1.85 (0.49)	<0.001	0.33 (0.53)	0.534	0.48 (0.49)	0.329
Black	-0.90 (0.50)	0.074	-2.79 (0.47)	<0.001	0.54 (0.52)	0.299	-0.03 (0.48)	0.950

PIAT=Peabody individual achievement test; AFQT=armed forces qualification test; HOME=home observation for measurement of the environment.

*Individual components of PIAT.

scores. None of them was significantly different from zero. In both cases, we also tested for any differences between the sibling pairs in the confounding factors used in the earlier analysis. None was significant (results not shown), suggesting that the method controls for all those factors and there is no need for further analysis. In short, the sibling pairs analysis concurs with the earlier analysis in offering no support for a cognitive advantage of breast feeding once confounding is taken into account.

Meta-analysis

Our search yielded 431 references, 73 of which reported original data for full term infants. In 19 reports the analysis explicitly controlled for parental IQ and in one further report¹⁰ it implicitly controlled for parental IQ via sibling comparisons. The 19 reports used data from 12 studies because studies that followed children prospectively and assessed them at different ages often repeated the analysis and published the results separately. When the published results did not present the data in a comparable format, we approached the corresponding author directly.

The figure summarises eight of the 12 studies,²⁷⁻³⁴ together with the results for PIAT total score from our study. We excluded four studies because the data were no longer available in a suit-

able form,³⁵ "nearly 100% had some breast feeding"³⁶ (T Vik, personal communication), or we could not contact the authors.^{37, 38} When a study assessed children at more than one age we chose the oldest age available as childhood measures of cognitive ability increase in reliability with age.³⁹ When more than one measure of ability was assessed we took the more general measure.

The studies with the biggest IQ advantage for breast feeding were those with smaller sample sizes and that controlled for fewer of the important additional confounders—namely, socioeconomic status, maternal education, maternal age, and HOME score. The study with the largest effect had a sample that was socially heterogeneous (D Hay, personal communication) and the next largest one that was "disadvantaged, consisting almost exclusively of members of social classes IV and V."³² Two studies assessed the children at only 2 years of age when the available measures are "best characterized as indices of mental development, and not IQ."³⁸

The greatest support for an effect of breast feeding came from the study by Fergusson et al,²⁷ which was based on the previous largest sample and from a birth cohort with an outstanding follow-up rate. It did not, however, control for maternal age,

Table 4 Supplementary analysis: omitting mother's IQ, and duration of breast feeding (only in those breast fed)

	PIAT-total		Maths*		Reading*		Comprehension*	
	B (SE)	P value	B (SE)	P value	B (SE)	P value	B (SE)	P value
Breast feeding†	1.42 (0.37)	<0.001	1.33 (0.35)	<0.001	1.14 (0.38)	0.002	1.32 (0.35)	<0.001
Duration of breast feeding								
BF (months)‡	0.11 (0.05)	0.025	0.07 (0.05)	0.127	0.08 (0.05)	0.102	0.14 (0.05)	0.003
Not breast fed	0.00		0.00		0.00		0.00	
1-5 weeks	0.32 (0.51)	0.526	0.04 (0.49)	0.934	0.33 (0.53)	0.533	0.47 (0.49)	0.334
6-12 weeks	0.18 (0.52)	0.734	0.32 (0.50)	0.514	0.01 (0.54)	0.980	0.21 (0.50)	0.681
13-28 weeks	0.65 (0.57)	0.254	0.86 (0.54)	0.112	0.44 (0.58)	0.449	0.24 (0.54)	0.661
≥29 weeks	1.50 (0.59)	0.011	1.41 (0.56)	0.012	1.00 (0.61)	0.099	1.72 (0.57)	0.002

PIAT=Peabody individual achievement test.

*Individual components of PIAT.

†Mutually adjusted model as in table 3, but omitting mother's AFQT score.

‡Duration of breast feeding (in months) as a continuous predictor; 1587 mothers, 2454 children, 7416 assessments.

Table 5 Supplementary analysis—sibling pairs

Breast feeding	PIAT-total		Maths*		Reading*		Comprehension*	
	Difference (SE)	P value	Difference (SE)	P value	Difference (SE)	P value	Difference (SE)	P value
Status†	-0.63 (0.94)	0.506	0.02 (1.01)	0.984	-1.00 (0.96)	0.296	-0.54 (1.00)	0.589
Duration‡	-0.13 (0.76)	0.866	-0.11 (0.82)	0.890	-0.41 (0.79)	0.602	0.20 (0.79)	0.796

PIAT=Peabody individual achievement test.

*Individual components of PIAT.

†Breastfed sibling minus non-breastfed sibling; n=332 pairs.

‡Sibling with longer duration minus sibling with shorter duration (n=545 pairs).

Research

studies combined provide no support for a beneficial effect of breast feeding.

Wider application of results

The generalisability of the results presented here must be considered carefully. This study and the others included in the meta-analysis are all based on samples from developed countries: United States,^{10 32 41} Europe,^{28 30 33} Australia,³⁴ and New Zealand.²⁷ Generalisation of the findings beyond these and similar societies would be unwise. We have also excluded premature and low birthweight infants for whom the effect may be different.

The broader context of breast feeding should also be considered. Evidence showing the many benefits of breast feeding for the child and mother led the World Health Organization and UNICEF to formulate the Innocenti Declaration,⁴² which includes exclusive breast feeding for 4-6 months as a global goal. Many of these benefits are equally applicable to developed countries.^{43 44} Even if it does not enhance intelligence, breast feeding remains "an unequalled way of providing ideal food for the healthy growth and development of infants."⁴⁵

We are grateful to Alex Ghys, Dale Hay, and Sandra Jacobson for providing data for the meta-analysis.

Contributors: The idea for the paper was generated during discussions between the authors. GD conducted the analysis, wrote the first draft of the manuscript, and is guarantor. All authors made substantial contributions to subsequent revisions.

Funding: GD is employed by the Medical Research Council. GDB is funded by a Wellcome advanced training fellowship and is also affiliated to the University of Edinburgh, Department of Psychology. ID is the recipient of a Royal Society-Wolfson research merit award.

Competing interests: None declared.

Ethical approval: Not required.

- 1 Hoefler C, Hardy M. Later development of breast fed and artificially fed infants. *JAMA* 1929;92:615-20.
- 2 Lucas A, Morley R, Cole TJ, Gore SM. A randomised multicentre study of human milk versus formula and later development in preterm infants. *Arch Dis Child Fetal Neonatal Ed* 1994;70:F141-6.
- 3 Lucas A, Morley R, Cole TJ, Lister G, Leeson-Payne C. Breast-milk and subsequent intelligence quotient in children born preterm. *Lancet* 1992;339:261-4.
- 4 Morley R, Cole TJ, Powell R, Lucas A. Mother's choice to provide breast milk and developmental outcome. *Arch Dis Child* 1988;63:1382-5.
- 5 Anderson JW, Johnstone BM, Remley DT. Breast-feeding and cognitive development: a meta-analysis. *Am J Clin Nutr* 1999;70:525-35.
- 6 Drane DL, Logemann JA. A critical evaluation of the evidence on the association between type of infant feeding and cognitive development. *Paediatr Perinat Epidemiol* 2000;14:349-56.
- 7 Jain A, Concato J, Leventhal JM. How good is the evidence linking breastfeeding and intelligence? *Pediatrics* 2002;109:1044-53.
- 8 Bouchard T, McGue M. Genetic and environmental influences on human psychological differences. *J Neurobiol* 2003;54:4-45.
- 9 Silva PA, Buckfield P, Spears GF. Some maternal and child developmental characteristics associated with breast feeding: a report from the Dunedin multidisciplinary child developmental study. *Aust Paediatr J* 1978;14:265-8.

What is known already on this topic

Breast feeding has many benefits for both child and mother

Breastfed children tend to score higher on intelligence tests

There are important differences between mothers who breast feed and those who do not

What this study adds

The apparent effect of breast feeding on intelligence in offspring is accounted for by characteristics of the mother and the home environment

The mother's own intelligence plays the largest part in this explanation

- 10 Evenhouse E, Reilly S. Improved estimates of the benefits of breastfeeding using sibling comparisons to reduce selection bias. *Health Services Res* 2005;40:1781-802.
- 11 Center for Human Resource Research. *NLSY79 users guide*. Columbus, OH: Ohio State University, 2004.
- 12 Center for Human Resource Research. *NLSY79 child and young adult data users guide*. Columbus, OH: Ohio State University, 2002.
- 13 Lucas A, Morley R, Cole TJ, Gore SM, Lucas PJ, Crowle P, et al. Early diet in preterm babies and developmental status at 18 months. *Lancet* 1990;335:1477-81.
- 14 Keck C. The initiation and duration of breastfeeding among employed women in the United States. Columbus, OH: Ohio State University, 1997 (PhD thesis).
- 15 Quo G, Harris KM. The mechanisms mediating the effects of poverty on children's intellectual development. *Demography* 2000;37:431-47.
- 16 Johnson DL, Swank PR, Howie VM, Baldwin CD, Owen M, Luttman D. Does HOME add to the prediction of child intelligence over and above SES? *J Genetic Psychol* 1993;154:33-40.
- 17 Pollock J. Long-term associations with infant feeding in a clinically advantaged population of babies. *Dev Med Child Neurol* 1994;36:429-40.
- 18 Rey J. Breastfeeding and cognitive development. *Acta Paediatr Suppl* 2003;92:11-8.
- 19 Caldwell BM, Bradley RH. *Home observation for measurement of the environment*. Little Rock, AR: University of Arkansas, 1984.
- 20 Diggle PJ, Liang KL, Zeger SL. *Analysis of longitudinal data*. Oxford: Oxford University Press, 1994.
- 21 Goldstein H. *Multilevel statistical models*. 2nd ed. London: Arnold, 1995.
- 22 Bauml ER, Carvajal S, Harrist RB. Analysis of repeated measures data. In: Reise SP, Duan N, ed. *Multilevel modeling: methodological advances, issues, and applications*. Mahwah, NJ: Lawrence Erlbaum, 2003:140-56.
- 23 Cnaan A, Laird NM, Slator P. Using the general linear mixed model to analyse unbalanced repeated measures and longitudinal data. *Stat Med* 1997;16:2349-80.
- 24 Singer JD. Using SAS PROC MIXED to fit multilevel models, hierarchical models, and individual growth models. *J Educ Behav Stat* 1998;23:323-55.
- 25 Willett JB, Singer JD, Martin NC. The design and analysis of longitudinal studies of development and psychopathology in context: statistical models and methodological recommendations. *Develop Psychopathol* 1998;10:395-426.
- 26 Hill AB. *Principles of medical statistics*. New York: Oxford University Press, 1971.
- 27 Fergusson DM, Beautrais AL, Silva PA. Breast-feeding and cognitive development in the first seven years of life. *Soc Sci Med* 1982;16:1705-8.
- 28 Ghys A, Bakker E, Hornstra G, van den Hout M. Red blood cell and plasma phospholipid arachidonic and docosahexaenoic acid levels at birth and cognitive development at 4 years of age. *Early Hum Dev* 2002;69:83-90.
- 29 Gomez-Sanchiz M, Canete R, Rodero I, Baeza JE, Gonzalez JA. Influence of breast-feeding and parental intelligence on cognitive development in the 24-month-old child. *Clin Pediatr (Phila)* 2004;43:753-61.
- 30 Hay DF, Pawby S, Sharp D, Asten P, Mills A, Kumar R. Intellectual problems shown by 11-year-old children whose mothers had postnatal depression. *J Child Psychol Psychiatry* 2001;42:871-89.
- 31 Jacobson SW, Chiodo LM, Jacobson JL. Breastfeeding effects on intelligence quotient in 4- and 11-year-old children. *Pediatrics* 1999;103:e71.
- 32 Morrow-Tlucak M, Haude RH, Ernhart CB. Breastfeeding and cognitive development in the first 2 years of life. *Soc Sci Med* 1988;26:635-9.
- 33 Richards M, Wadsworth M, Rahimi-Foroushani A, Hardy R, Kuh D, Paul A. Infant nutrition and cognitive development in the first offspring of a national UK birth cohort. *Dev Med Child Neurol* 1998;40:163-7.
- 34 Wigg NR, Tong SL, McMichael AJ, Baghurst PA, Vimpani G, Roberts R. Does breastfeeding at six months predict cognitive development? *Austr N Z J Public Health* 1998;22:232-6.
- 35 Johnson DL, Swank PR, Howie VM, Baldwin CD, Owen M. Breast feeding and children's intelligence. *Psychol Rep* 1996;79:1179-85.
- 36 Angelsen NK, Vik T, Jacobsen G, Bakketeig LS. Breast feeding and cognitive development at age 1 and 5 years. *Arch Dis Child* 2001;85:183-8.
- 37 de Andraca I, Pino P, de la Parra A, Rivera F, Castillo M. [Risk factors for psychomotor development among infants born under optimal biological conditions]. *Rev Saude Publica* 1998;32:138-47.
- 38 Vreugdenhil HJ, Mulder PG, Emmen HH, Weisglas-Kuperus N. Effects of perinatal exposure to PCBs on neuropsychological functions in the Rotterdam cohort at 9 years of age. *Neuropsychology* 2004;18:185-93.
- 39 Wilson RS. The Louisville twin study: developmental synchronies in behavior. *Child Dev* 1983;54:298-316.
- 40 Thompson SG, Higgins JPT. How should meta-regression analyses be undertaken and interpreted? *Stat Med* 2002;21:1559-73.
- 41 Jacobson S, Jacobson J. Breastfeeding and intelligence. *Lancet* 1992;339:926.
- 42 WHO/UNICEF. *The Innocenti declaration on the protection, promotion and support of breastfeeding*. Geneva: WHO/UNICEF, 1990.
- 43 American Academy of Pediatrics. Policy statement. Breastfeeding and the use of human milk. *Pediatrics* 1997;100:1035-9.
- 44 Yngve A, Sjostrom M. Breastfeeding in countries of the European Union and EFTA: current and proposed recommendations, rationale, prevalence, duration and trends. *Public Health Nutr* 2001;4:631-45.
- 45 World Health Organization. *Global strategy for infant and young child feeding*. Geneva: WHO, 2000.

(Accepted 30 August 2006)

doi: 10.1136/bmj.38978.699583.55

MRC Social and Public Health Sciences Unit, Glasgow G12 8RZ

Geoff Der *statistician*

G David Batty *Wellcome fellow*

Department of Psychology, University of Edinburgh, Edinburgh

Ian J Deary *professor of differential psychology*

Correspondence to: G Der Geoff@msoc.mrc.gla.ac.uk