

Efficacy of a smoking cessation intervention using the AHCPR guideline tailored for Koreans: a randomized controlled trial

JANG-RAK KIM¹, MYOUNG-SOON LEE², JIN-YONG HWANG³ and JONG DEOG LEE³

¹Department of Preventive Medicine, College of Medicine, and Institute of Health Sciences, Gyeongsang National University, Jinju, South Korea, ²Department of Social Medicine, School of Medicine, Sung Kyun Kwan University, Suwon, South Korea and ³Department of Internal Medicine, College of Medicine, Gyeongsang National University, Jinju, South Korea

ABSTRACT

This study was undertaken to evaluate a tailored smoking cessation intervention, which is applicable to Korean culture, using the Agency for Health Care Policy and Research (AHCPR) guideline. On-site counselors provided brief nurse-assisted smoking cessation counseling, including follow-up telephone support, to prevent a relapse in 200 randomly assigned smoking patients. These patients were referred by their physicians regardless of their willingness in smoking cessation in the outpatient department at a university hospital. Nicotine replacement therapy was not provided. Another 201 patients served as a control. After 5 months, current smoking cessation was

self-reported on the phone and validated later by a portable carbon monoxide analyzer. After 5 months, the participants in the intervention group were no more likely to quit smoking than the control group. A subgroup analysis by age showed that the intervention among 166 younger smokers (aged 49 or less) was significantly more likely to be effective [risk ratio = 5.76 [95% confidence interval (CI) 1.34–24.74]] than it was among 235 older smokers (aged 50 or more) [risk ratio = 1.03 (95% CI 0.53–1.99)]. This study suggests a smoking cessation intervention using the AHCPR guideline tailored for Koreans, is effective for assisting outpatients aged 49 or younger to quit smoking.

Key words: Korean; randomized controlled trial; smoking cessation

INTRODUCTION

Cigarette smoking is the most important preventable cause of morbidity and mortality in society (US Department of Health and Human Services, 1989). Tobacco accounted for 24% of all male deaths and 7% of all female deaths in developed countries in 1990 (Peto *et al.*, 1996) and for 17% of the total number of deaths in Korean men in 1985 (Meng, 1988). Extensive evidence on the health benefits of smoking cessation has accumulated. With few exceptions, disease risks are reduced following smoking

cessation and continue to fall as abstinence is maintained (Samet, 1992).

Clinicians should take an active role in reducing the prevalence of smoking through effective smoking cessation intervention. Unfortunately, many are unaware of how to treat smoking patients because they are unacquainted with the available, effective and brief interventions for clinical settings. The Agency for Health Care Policy and Research (AHCPR, presently, the Agency for Healthcare Research and Quality,

AHRQ) developed and published a clinical guideline for smoking cessation in 1996 after reviewing approximately 3000 articles (Fiore *et al.*, 1996). In 2000, a consortium of governmental and non-profit organizations including the AHRQ updated the guideline. However, there were no major differences between the original and the revised one. The AHCPR guideline states the major steps (the '5 As') to intervention in the primary care setting are: (1) ask the patient if he or she uses tobacco; (2) advise him or her to quit; (3) assess the patient's willingness to quit; (4) assist those who are willing to quit; and (5) arrange for follow-up contact to prevent relapse (The Tobacco Use and Dependence Clinical Practice Guideline Panel, Staff, and Consortium Representatives, 2000).

As health behaviors such as smoking are composite expressions of social and cultural circumstances (Green and Kreuter, 1991), interventions developed in western countries may not be applicable or effective in other countries. Although few studies have examined the AHCPR guideline tailored to a particular ethnic group, some types of tailoring such as the use of appropriate language and messages can increase treatment effectiveness (Fiore *et al.*, 1996; King *et al.*, 1997).

In Korea, the prevalence of smoking was quite high (61.8%) for men but lower (5.4%) for women in the year 2001 (Choi and Kim, 2002). Most Korean physicians believe that smoking cessation intervention is important to assist smoking patients to quit. However, they lack training in the major steps (the '5 As') from the AHCPR guideline (Lee *et al.*, 1991). Additionally, the most preferred intervention is only to advise the patients to quit smoking with an explanation of the health risks involved (Yi *et al.*, 1995). Still, the AHCPR guideline has not been tried in clinical settings in Korea, nor translated into Korean.

We performed this study to determine the effectiveness of the AHCPR smoking cessation intervention guideline, in particular, when tailored for a Korean population.

METHODS

Study subjects

The study participants were patients who visited their doctors in the outpatient department of internal medicine at Gyeongsang National

University Hospital between January and March, 2000. The inclusion criteria were: (1) currently smoking one cigarette a day or more on average; (2) age ≥ 20 years; (3) providing oral informed consent; (4) having no terminal stage illness; (5) willing to participate in the study; (6) providing a telephone number and (7) able to understand the interview.

A yellow sheet inserted into every medical chart reminded all physicians to ask their patients about their current smoking status (as the strategy of 'Ask' according to the AHCPR guideline). According to the study protocol, the physicians were requested to provide routinely concise and strong stop-smoking advice to all smoking patients (as the strategy of 'Advise' according to the AHCPR guideline), and to encourage them to visit one of the three on-site counselors' desks in the lobby. The on-site counselors were retired nurses who were trained for this study. 401 people agreed to participate in this study among the 408 smoking patients who visited the on-site counselors. Of these, 394 (98.3%) could be followed up for at least 5 months.

We randomly assigned the study participants to either the intervention or the control group. This was consummated according to a random list determined by fixed randomization (Meinert and Tonascia, 1986) with an allocation ratio of 1:1, a block size of 6 and 12 allocation strata according to their willingness to quit within 1 month or not, whether they were scheduled to be admitted within 1 month or not, and which of the three on-site counselors they saw. The treatment assignments based on each of the 12 stratum were placed in sealed opaque envelopes which the counselors opened at the formal enrollment of the study participants.

A flowchart of the study design is provided in Figure 1.

Baseline assessment

After obtaining informed consent, the on-site counselors gathered baseline information using a structured interview. The information regarding potential confounders included: (1) whether they were willing to quit smoking in 1 month (as the strategy of 'Assess' according to the updated AHCPR guideline), (those who were willing to quit smoking within 1 month were referred to as being in the preparation stage of change) (DiClemente *et al.*, 1991). (2) Whether they were




Fig. 1: Flowchart of the study design. * indicates that the 12 strata were based on whether or not participants were willing to quit within 1 month, whether or not they were scheduled to be admitted within 1 month, and which of the 3 on-site counselors they saw.

scheduled to be admitted within 1 month. (3) Socio-demographic factors such as age, sex, educational level and occupation. (4) Whether they lived together with spouses, smokers or children. (5) Whether or not they attempted to seriously quit for >1 day. (6) What illnesses

they had. (7) A Fagerström test for nicotine dependence (FTND). (8) Self-efficacy to avoid smoking.

A high willingness to quit smoking, hospitalization, the presence of a non-smoking spouse, other smokers, or a child in the home, as well as

previous quit attempts are related to smoking cessation (Fiore *et al.*, 1996; Dale *et al.*, 1997; Okah *et al.*, 2002).

The serious illnesses associated with smoking are heart diseases such as ischemic heart disease, hypertension and arrhythmia, lung diseases such as chronic bronchitis, emphysema and asthma, or malignant tumors. It is believed that patients with such diseases were more motivated to quit smoking as reported in another study (Wilkes and Evans, 1999).

The FTND is a revision of the Fagerström tolerance questionnaire (FTQ), and is a predictor of smoking cessation (Kozlowski *et al.*, 1994). The FTND consists of six questions; 'How soon after you wake up do you smoke your first cigarette?', 'Do you find it difficult to refrain from smoking in places where it is forbidden?', 'Which cigarette would you hate most to give up?', 'How many cigarettes/day do you smoke?', 'Do you smoke more frequently during the first hours after waking than during the rest of the day?' and 'Do you smoke even if you are so ill that you are in bed most of the day?'. The FTND scores range from 0 to 10 (Heatherton *et al.*, 1991).

Self-efficacy is a predictor of relapse over time (Gulliver *et al.*, 1995). It was measured by a four-item assessment of the patients' confidence to abstain from smoking in high-risk situations: 'when they are feeling depressed', 'when they are on vacation and want to relax', 'when they are feeling a craving for a cigarette' and 'when they are being offered a cigarette in a social situation'. The self-efficacy score was derived by averaging the Likert-scale answers (1 = not at all, 2 = not very, 3 = moderately, 4 = very and 5 = extremely) across the four items (Coon *et al.*, 1998). In the few studies investigating Korean smokers, age, educational level, smoking amount, FTND scores and the presence of a serious illnesses associated with smoking were related to successful smoking cessation in outpatients (Won *et al.*, 1992; Kim *et al.*, 1995; Cho *et al.*, 1996; Kim and Seo, 2001).

Interventions

For 132 of 200 participants in the intervention group, who were willing to quit smoking within 1 month, the intervention given by on-site counselors after baseline assessment was 2A of the AHCPR guideline, i.e. to *assist* and to *arrange* (follow-up contact).

The specific intervention actions were: helping participants set a quit date, arrange signing of a

stop-smoking contract, and providing self-help material with brief explanations addressing the key advice on successful quitting under the AHCPR guideline. The culturally specific intervention tailorings were: to use language appropriate to Koreans, to supply a plastic cigarette-sized tube as a cigarette substitute (according to a preliminary survey showing that many Korean smokers in the study area had smoked to pass the time) and to have a picture (particularly from the grandchildren) on the cover of self-help material addressing the importance of family support, in smoking cessation.

Information on nicotine replacement therapy was not given in order to evaluate the efficacy of behavioral intervention alone. The average time needed to supply all intervention steps was 11.1 ± 3.8 min.

Additional interventional actions included mailing postcards to remind the patients of the quit date, and counseling via telephone 1 week and 1 month after the quit date. The counselors on the phone congratulated any successful quit attempts, or reviewed and elicited a recommitment to total abstinence if smoking had occurred. Overall, the interview took approximately 7 min per call.

For the remaining 68 participants in the intervention group who were unwilling to quit within 1 month, the intervention provided by on-site counselors was in the form of the 4Rs (relevance, risks, rewards and repetition) according to the AHCPR guideline. This was provided initially on-site and then over the phone after 2–3 weeks, and again after 6–7 weeks to try to motivate them to quit smoking.

The 201 participants in the control group were told by on-site counselors to quit smoking by their own free will without any further assistance.

A summary of the randomization and intervention scheme is provided in Figure 1.

The institutional review board of Gyeongsang University Hospital approved this study.

Outcome assessment

The self-reported serious quit attempts (defined as temporary smoking abstinence for at least 1 day) and smoking cessation (defined as absolutely no smoking since the last quit attempt) were assessed by a telephone interview at least 5 months after the participants in both the intervention and control group were enrolled in the study. Additionally, participants were informed that biochemical samples might be

taken from those who stated that they had quit. Five months was chosen to examine only clinically important outcomes (i.e. long-term cessation) according to the AHCPR guideline for smoking cessation (Fiore *et al.*, 1996).

The interviewers were on-site counselors, and were unaware of the participants' group assignment. The self-reported current smoking cessation was validated at home or at the outpatient department within 3 weeks at the latest using the carbon monoxide (CO) cut-off values of 7 ppm or less (Kawane and Soejima, 1992) in the expired alveolar air from a deep breath by a portable CO analyzer (Smokerlyzer®, Bedfont Scientific). However, the half-life of CO is only 3–5 h (Windsor and Orleans, 1986). Therefore, even if the alveolar CO level was under 7 ppm, the interviewees who confessed smoking even a puff or more since their last quit date were classified as smokers (six persons in the intervention group, four persons in the control group).

Statistical analysis

Those who refused to take alveolar CO analysis (eight persons in the intervention group, three persons in the control group) among the self-reported quitters or could not be followed up (four persons in the intervention group, three persons in the control group) were also classified as smokers. This complies with the intention-to-treat analysis, which preserves the full value

of randomization, and guarantees the control over the baseline confounders (Hulley and Cummings, 1988).

The efficacy of the intervention was determined by comparing outcome measures between groups. The risk ratio (the ratio of the proportions of the intervention to the control group that had successfully quit smoking) was calculated with a 95% confidence interval (CI) in all participants. Subgroup analysis based on the participants' willingness to quit within 1 month, scheduled admission within 1 month, and age was planned a priori to determine which subgroup could benefit most from the intervention. Multivariate analysis using multiple logistic regression was done to adjust the efficacy for the baseline potential confounders shown in Table 1. All the data were analyzed by SPSS for Windows (version 7.5) (SPSS Inc., Chicago, IL).

RESULTS

The two groups were similar in terms of the proportions of those who were willing to quit smoking within 1 month and of those who were to be admitted within 1 month, which were used as the stratifying variables. There were no statistically significant differences in the potential confounders such as age, sex, educational level, occupation, whether they lived together with smokers, children or spouses, smoking behavior

Table 1: Baseline characteristics of the study participants

Characteristics	Intervention group (<i>n</i> = 200)	Control group (<i>n</i> = 201)
Willingness to quit within 1 month (%)	132 (66.0)	129 (64.2)
Scheduled admission within 1 month (%)	13 (6.5)	13 (6.5)
Age (years) (mean ± SD)	51.6 ± 13.0	53.1 ± 13.5
Men (%)	188 (94.5)	180 (90.5)
High school graduation (%)	108 (54.0)	102 (50.7)
Farming (%)	47 (23.5)	57 (28.4)
Serious illness ^{a,b} (%)	83 (41.5)	105 (52.2)
Living together with smokers (%)	49 (24.5)	47 (23.4)
Living together with children or babies (%)	44 (22.0)	38 (18.9)
Living together with spouse (%)	177 (88.5)	170 (84.6)
Previous serious quit attempt(s) (%)	148 (74.0)	147 (73.1)
FTND ^c (mean ± SD)	4.0 ± 2.2	4.1 ± 2.2
Self-efficacy ^d (mean ± SD)	3.7 ± 0.9	3.7 ± 1.0

^a*p* < 0.05 by χ^2 or *t*-test.

^bHeart diseases such as ischemic heart disease, hypertension and arrhythmia, lung diseases such as chronic bronchitis, emphysema and asthma, or malignant tumor.

^cFagerström test for nicotine dependence (scores range from 0 to 10).

^dSelf-efficacy to avoid cigarette smoking is the average of Likert-scale answers (1 = not at all, 2 = not very, 3 = moderately, 4 = very and 5 = extremely) to the four high-risk situations.

and self-efficacy to avoid smoking between the two groups. The only exception was for the serious illness confounder, where more participants in the control group had serious illnesses (Table 1).

During the 5 months follow-up stage, participants in the intervention group were 1.20 times (95% CI 1.04–1.40) more likely to report serious quit attempts than the control group [69.5 (139/200) versus 57.7% (116/201)]. After 5 months, participants in the intervention group were 1.86 times (95% CI 1.20–2.87) more likely to report smoking cessation than the control group [24.0 (48/200) versus 12.9% (26/201)]. However, the validated quit rate after 5 months was no more likely to be higher in the intervention group than in the control group (Table 2).

Subgroup analysis by age revealed that the validated quit rate after 5 months was 5.76 times (95% CI 1.34–24.74) higher in the intervention group than in the control group [14.8 (13/88) versus 2.6% (2/78)] among 166 participants who were 49 years or younger; while the rates were similar in the two groups among 235 participants who were 50 years or older (Table 3).

The validated quit rate ratios after 5 months in the intervention and control group did not depend on whether or not the participants were willing to quit within 1 month, or whether they

were to be admitted within 1 month or not in other subgroup analyses (data not shown).

Multivariate analysis to adjust the efficacy for the baseline potential confounders shown in Table 1 made little difference. The crude odds ratio, 1.66 (95% CI 0.88–3.10), for the efficacy of the intervention was changed to 1.70 (95% CI 0.84–3.41) in the multiple logistic regression (data not shown).

DISCUSSION

As a minimum standard, smoking cessation research must combine self-reports with an objective verification (Windsor and Orleans, 1986). Much effort was made to confirm self-reported abstinence. The interviewers were blinded to the participants' group assignment. They told participants that biochemical samples might be taken from quitters, prior to asking questions regarding their smoking abstinence on the telephone. Verification of the self-report was made by a portable CO analyzer (Smokerlyzer®, Bedfont Scientific) using the CO cut-off value of 7 ppm or less. This method was inexpensive and required minimal staff training. Although the half-life of CO is only 3–5 h, and is influenced by

Table 2: Number, proportion (%), and risk ratios of serious quit attempts, during the 5 months, self-reported quits after 5 months, and validated quits after 5 months

	Intervention group (<i>n</i> = 200)	Control group (<i>n</i> = 201)	Risk ratio ^a (95% CI ^b)
Serious quit attempts during 5 months	139 (69.5)	116 (57.7)	1.20 (1.04–1.40)
Self-reported quits after 5 months	48 (24.0)	26 (12.9)	1.86 (1.20–2.87)
Validated quits after 5 months	28 (14.0)	18 (9.0)	1.56 (0.89–2.73)

^aRatio of the proportions (%) of the intervention to the control group.

^bCI, confidence interval.

Table 3: Number, proportion (%), and risk ratios of validated quits after 5 months in subgroup analysis based on age

	Intervention group	Control group	Risk ratio ^a (95% CI ^b)
Age <50 years	<i>n</i> = 88	<i>n</i> = 78	
Validated quits after 5 months	13 (14.8)	2 (2.6)	5.76 (1.34–24.74)
Age ≥50 years	<i>n</i> = 112	<i>n</i> = 123	
Validated quits after 5 months	15 (13.4)	16 (13.0)	1.03 (0.53–1.99)

^aRatio of the proportions (%) of the intervention to the control group.

^bCI, confidence interval.

multiple non-smoking-related CO sources, some studies have found it useful to distinguish between smokers and non-smokers, and to detect changes in the smoking rate (Windsor and Orleans, 1986).

Only 28 (58.3%) of the 48 self-reported abstinences in the intervention group and 18 (69.2%) of 26 self-reported abstinences in the control group were confirmed. The reason why more self-reports were not reliable in the intervention group could be that more participants in the intervention group did not wish to disappoint the on-site counselors with their failures in smoking cessation in spite of the assistance.

For the intention-to-treat analysis, all participants lost to follow-up were classified as smokers. We could follow up 394 (98.3%) of the 401 participants for at least 5 months after enrollment. The high follow-up rate was partly due to the fact that many participants had cellular phones.

The intervention in this study, a tailored AHCPR guideline of the '5 As' strategies (ask, advise, assess, assist, arrange), was applied in an outpatient setting in Korea with no problems.

This brief intervention increased both the serious attempts to quit smoking during the 5 months follow-up period and the validated quit rate after 5 months. Although not statistically significant, the ratio of the validated quit rate in the intervention group to that in the control group, 1.56 (1.66 in terms of odds ratio) is similar to the odds ratio of 1.4 (95% CI 1.1–1.8) of the interventions delivered by non-physician medical health care providers in a meta-analysis of 20 studies (Fiore *et al.*, 1996). This suggests that smoking cessation interventions developed in western countries, with a little tailoring, may be applied and effective in other countries such as Korea, which have quite different social and cultural circumstances.

Current smokers can be grouped in each of the stages of change: the pre-contemplation stage of not seriously considering quitting within the next 6 months, the contemplation stage of seriously considering quitting within the next 6 months, and the preparation stage of planning to quit within 1 month. Smoking cessation intervention can increase successful quitting rates by being sensitive to the patient's stage and by shifting strategies according to the stage of change (DiClemente *et al.*, 1991). The assessment of the stages (as a strategy of 'Assess') in this study showed that 384 (95.8%) participants wanted to quit smoking, 261 (65.1%) were in the preparation

stage and 71 (17.7%) were in the contemplation stage. The validated quit rate (14.9%) of those who were in the preparation stage was much higher than of those (5.0%) of other stages (data not shown).

In the pilot study, the number of new patients was estimated to be 2700 for 3 months, and the smoking prevalence among them was 22%. The low smoking prevalence in female patients and the presence of many ex-smokers (some may have been lying to their physicians when they stated that they quit smoking) contributed to this overall low prevalence. Therefore, the estimated number of smoking patients was 594, which means the sample size of this study, 401, covered 67.5% of the smoking patients. More motivated patients might have participated in our study.

For the 68 participants in the intervention group who were not willing to quit within 1 month, the intervention provided was the 4R messages (relevance, risks, rewards, repetition) in order to motivate them to quit smoking on-site and on the phone after 2–3 weeks, and again after 6–7 weeks. Fourteen (20.6%) participants expressed their determination to try to quit during this process, and were later given the same intervention as the motivated participants in the intervention group. However, the validated quit rate in the intervention group was only 5.9 (4/68) compared with 4.2% (3/72) in the control group. The efficacy of motivational intervention is still unknown (Fiore *et al.*, 1996).

The three stratifying variables (a participant's willingness to quit, scheduled admission and age) were pre-specified before analysis, because there may have been problems of multiple hypotheses in subgroup analyses (Hulley and Cummings, 1988). The efficacy results did not depend on the participants' willingness to quit or their scheduled admission. However, age made a big difference. The participants in the intervention group were 5.76 times (95% CI 1.34–24.74) more likely to quit smoking than the control group if they were 49 years or younger; while the quit rates were similar in the two groups for those participants who were 50 years or older. Younger participants in this study were much more educated than the older ones. They may have been better informed and more receptive to new ideas than older people in a rapidly developing country like Korea. However, the risk ratio in our study had a very wide confidence interval because of small sample sizes in the subgroups. To get a precise ratio, additional studies with

more participants are needed. Other studies have also suggested lower efficacy results and the need for better tailored smoking cessation interventions for older smokers (Blurton *et al.*, 1995; Morgan *et al.*, 1996; Ioka *et al.*, 2001).

One of the critical reasons why Korean doctors do not deliver smoking cessation counseling is a 'lack of time' (Lee *et al.*, 1991). Physicians can fulfil their responsibility by designing a system that delegates at least part of the preventive services to other personnel, such as nurses, while maintaining their ongoing involvement (Kottke *et al.*, 1993). Moreover, interventions delivered by nurses are much cheaper than those by doctors.

In conclusion, this study suggests that smoking cessation intervention using the AHCPR guideline tailored for Koreans can be easily implemented in an outpatient setting and is effective for outpatients who are 49 years or younger. However, more interventional studies, in order to obtain more reliable information in Korea, should be implemented.

ACKNOWLEDGEMENTS

This work was supported by a grant from the Korean Health Promotion Fund (KIHSA HC-03-17). Special thanks are given to the nurses and physicians working at the outpatient department of internal medicine at Gyeongsang National University Hospital for participating in our project.

Address for correspondence:

Jang-Rak Kim
Department of Preventive Medicine
College of Medicine
Gyeongsang National University
92 Chilam-Dong
Jinju, 660-751
South Korea
E-mail: jrkim@nongae.gsnu.ac.kr

REFERENCES

- Blurton, L. C., Paglia, M. J., German, P. S., Shapiro, S. and Damiano, A. M. (1995) The effect among older persons of general preventive visit on three health behaviors: smoking, excessive alcohol drinking, and sedentary lifestyle. The Medicare Preventive Services Research Team. *Preventative Medicine*, **24**, 492–497.
- Cho, H. J., Yeo, J. K. and Sunwoo, S. (1996) Predicting factors of smoking cessation in patients visiting smoking cessation clinic. *Journal of Korean Academy of Family Medicine*, **17**, 1211 (in Korean).
- Choi, J. S. and Kim, C. I. (2002) *2001 National Health and Nutrition Survey in Korea (Overview)*. Ministry of Health and Welfare, Korean Institute of Health and Social Affairs. Publication No. 2002-61-1 (in Korean).
- Coon, G. M., Pena, D. and Illich, P. A. (1998) Self-efficacy and substance abuse; assessment using a brief phone interview. *Journal of Substance Abuse Treatment*, **15**, 385–391.
- Dale, L. C., Olsen, D. A., Patten, C. A., Schroeder, D. R., Croghan, I. T., Hurt, R. D., Offord, K. P. and Wolter, T. D. (1997) Predictors of smoking cessation among elderly smokers treated for nicotine dependence. *Tobacco Control*, **6**, 181–187.
- DiClemente, C. C., Prochaska, J. O., Fairhurst, S. K., Velicer, W. F., Velasquez, M. M. and Rossi, J. S. (1991) The process of smoking cessation: an analysis of precontemplation, contemplation, and preparation stages of change. *Journal of Consulting and Clinical Psychology*, **59**, 295–304.
- Fiore, M. C., Bailey, W. C., Cohen, S. J., Dorfman, S. F., Goldstein, M. G., Gritz, E. R., *et al.* (1996) *Smoking Cessation. Clinical Practice Guideline No 18*. Rockville (MD), US Department of Health and Human Services, Public Health Service, Agency for Health Care Policy and Research. AHCPR Publication No. 96-0692.
- Green, L. W. and Kreuter, M. W. (1991) *Health Promotion Planning: An Educational and Environmental Approach*. Mayfield Publishing Company, Mountain View.
- Gulliver, S. B., Hughes, J. R., Solomon, L. J. and Dey, A. N. (1995) An investigation of self-efficacy, partner support and daily stresses as predictors of relapse to smoking in self-quitters. *Addiction*, **90**, 767–772.
- Heatherton, T. F., Kozlowski, L. T., Frecker, R. C., Fagerström, K. O. (1991) The Fagerström test for nicotine dependence: a revision of the Fagerström tolerance questionnaire. *British Journal of Addiction*, **86**, 1119–1127.
- Hulley, S. B. and Cummings, S. R. (1988) *Designing Clinical Research: An Epidemiologic Approach*. Williams & Wilkins, Baltimore, MD.
- Ioka, A., Nakamura, M., Shirokawa, N., Kinoshita, T., Masui, S., Imai, K., *et al.* (2001) Natural killer activity and its changes among participants in a smoking cessation intervention program—a prospective pilot study of 6 months' duration. *Journal of Epidemiology*, **11**, 238–242.
- Kawane, H. and Soejima, R. (1992) *Smoking habits and carbon monoxide concentration in expired air*. Proceedings of the 12th Asia Pacific congress on diseases of the chest, Seoul.
- Kim, C. H. and Seo, H. G. (2001) The factors associated with success of smoking cessation at smoking-cessation clinic. *Journal of Korean Academy of Family Medicine*, **22**, 1603–1611 (in Korean).
- Kim, M. K., Jin, C. K., Kim, Y. J. and Cho, B. M. (1995) The disease related factors affecting cessation of smoking. *J Korean Acad Fam Med*, **16**, 880–890 (in Korean).
- King, T. K., Borrelli, B., Black, C., Pinto, B. M., Marcus, B. H. (1997) Minority women and tobacco: implications for smoking cessation interventions. *Annals of Behavioral Medicine*, **19**, 301–313.
- Kottke, T. E., Brekke, M. L., Solberg, L. I. (1993) Making 'time' for preventive services. *Mayo Clinic Proceedings*, **68**, 785–791.
- Kozlowski, L. T., Porter, C. Q., Orleans, C. T., Pope, M. A. and Heatherton, T. (1994) Predicting smoking cessation with self-reported measures of nicotine

- dependence: FTQ, FTND, and HIS. *Drug Alcohol Dependency*, **34**, 211–216.
- Lee, K. S., Kim, H. S., Yang, S. J., Shin, H. C., Park, E. S., Lee, T. J., *et al.* (1991) Physicians' knowledge, attitude, and practice for quit smoking counseling. *Family Physician*, **12**, 35–43 (in Korean).
- Meinert, C. L. and Tonascia, S. (1986) *Clinical Trials*. Oxford University Press, New York, NY.
- Meng, K. H. (1988) Smoking-attributable mortality among Korean adults. *Korean Journal of Epidemiology*, **10**, 138–145 (in Korean).
- Morgan, G. D., Noll, E. L., Orleans, C. T., Rimer, B. K., Amfoh, K. and Bonney, G. (1996) Reaching midlife and older smokers: tailored interventions for routine medical care. *Preventative Medicine*, **25**, 346–354.
- Okah, F. A., Choi, W. S., Okuyemi, K. S. and Ahluwalia, J. S. (2002) Effect of children on home restriction by inner-city smokers. *Pediatrics*, **109**, 244–249
- Peto, R., Lopez, A. D., Boreham, J., Thun, M., Heath, C. Jr and Doll, R. (1996) Mortality from smoking worldwide. *British Medical Bulletin*, **52**, 12–21.
- Samet, J. M. (1992) The health benefits of smoking cessation. *Medical Clinics of North America*, **76**, 399–414.
- The Tobacco Use and Dependence Clinical Practice Guideline Panel, Staff, and Consortium Representatives (2000) A clinical practice guideline for treating tobacco use dependence. A US Public Health Service Report. *Journal of the American Medical Association*, **283**, 3244–3254.
- US Department of Health and Human Services. (1989) *Reducing the health consequences of smoking: 25 years of progress. A report of the Surgeon General*. Atlanta(GA), US Department of Health and Services, Public Health Service, Centers for Disease Control, Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health. DHHS Publication No. (CDC)89-8411.
- Wilkes, S. and Evans, A. A. (1999) A cross-sectional study comparing the motivation for smoking cessation in apparently healthy patients who smoke to those who smoke and have ischaemic heart disease, hypertension or diabetes. *Family Practitioner*, **16**, 608–610
- Windsor, R. A. and Orleans, C. T. (1986) Guideline and methodological standards for smoking cessation intervention research among pregnant women: improving the science and art. *Health Education Quarterly*, **13**, 131–161.
- Won, Y. I., Jeon, T. H., Lee, D. S., Oh, S. W., Choi, G. H., Yoo, T., *et al.* (1992) Baseline factors associated with smoking cessation. *Journal of Korean Academy of Family Medicine*, **13**, 862–868 (in Korean).
- Yi, S. H., Ha, Y. H. and Choi, H. R. (1995) The attitude to patient's smoking of the family physicians. *Journal of Korean Academy of Family Medicine*, **16**, 38–48 (in Korean).