

Influence of socio-economic deprivation on the prevalence and outcome of depression in primary care

The Hampshire Depression Project

K. OSTLER, C. THOMPSON, A.-L. K. KINMONTH, R. C. PEVELER, L. STEVENS
and A. STEVENS

Background Health inequalities exist for many disorders, but the contribution of deprivation to the prevalence and outcome of depressive symptoms in primary care populations has been infrequently studied.

Aims To examine the influence of Jarman under-privileged area (UPA) scores on the prevalence and outcome of depressive symptoms in general practice patients.

Method 18 414 patients attending 55 representative practices completed the Hospital Anxiety and Depression Scale and a questionnaire for employment status. Outcome of those screening positive was assessed at 6 weeks and 6 months.

Results The UPA score accounted for 48.3% of the variance between practices in prevalence of depressive symptoms. Attending a high UPA score practice predicted persistence of depressive symptoms to 6 months.

Conclusions The socio-economic deprivation of a practice locality is a powerful predictor of the prevalence and persistence of depressive symptoms.

Declaration of interest The study was funded by the Medical Research Council. No conflict of interest known.

Socio-economic deprivation, as measured by the Jarman under-privileged area (UPA) score (Jarman, 1983) and other measures, is associated with a high prevalence of disease, including severe mental illness (Thornicroft *et al*, 1992; Acheson, 1998). In the general population the prevalence of depression correlates with area deprivation (Eachus *et al*, 1996), individual characteristics such as unemployment (Meltzer *et al*, 1995) and employment grade (Stansfield & Marmot, 1992). However, consecutive consulting samples in primary care are potentially more useful to the clinician, reflecting the effects on day-to-day practice. Only the early study of Shepherd *et al* (1966) has taken such a primary care perspective and none has evaluated the relative influence of locality *v.* individual deprivation, nor the influence of deprivation on outcome. An opportunity to do so arose in the context of the Hampshire Depression Project (HDP), a randomised controlled trial of an educational package on depression (Thompson *et al*, 2000).

METHOD

Subjects

Fifty-five out of 224 (26%) practices in Hampshire, including 152 practitioners (69% of all partners), agreed to take part in a randomised controlled trial of an educational intervention. The practices were representative of Hampshire (see Table 1 in Thompson *et al*, 2000 for details). In four separate phases, from November 1994 to March 1997, consecutive adult patients (16 years and over) attending routine surgeries were screened by researchers distributing questionnaires in the waiting room. In each phase, screening continued until 50 (where there was one practitioner taking part) or 30 (where there was more than one study doctor in the practice) valid questionnaires per practitioner

had been obtained. Patients were excluded if they were too physically ill or unable to complete the questionnaire.

A total of 20 832 valid screening questionnaires were collected over the four screening phases. Duplicates were avoided by including only the first attendance of patients with an identical gender, date of birth and practice, and excluding forms where this was unavailable, leaving 18 414 patients (85.4% of the whole sample).

Measures of depression and employment

Two questionnaires were administered to each patient containing: (a) a brief explanation of the study, a request for demographic information and employment status in eight categories; (b) the Hospital Anxiety and Depression (HAD) scale (Zigmond & Snaith, 1983), which yields separate scores for symptoms of anxiety (A) and depression (D). Each scale has seven items, scores 0–3. There are no items related to biological symptoms of anxiety or depression, thus allowing exclusion of artificially high scores due to comorbidity with physical illness. The scale has been validated as a screening tool in general practice (Wilkinson & Barczak, 1988) and has been shown to be sensitive to change (Herrman, 1997).

There are two levels of case definition for the D scale: ‘possible depression’ at ≥ 8 and ‘probable depression’ at ≥ 11 . Both thresholds were used in this study. All patients with possible depression were sent another HAD by post 6 weeks and 6 months later. Patients who did not respond were sent another 2 weeks later. Questionnaires returned more than 12 weeks after the census date were excluded. At both of these times ‘remission’ was defined as a D score less than 8 and ‘improvement’ was defined as a score 50% or less than at the index consultation. For practice prevalence estimates, the higher threshold of ≥ 11 was used (probable depression), as it has a greater specificity against clinically diagnosed depression (Upadhyaya & Stanley, 1993).

At each consultation, practitioners rated depression on a 4-point scale. Ratings of 2: ‘clinically significant depressive illness, mild’, and 3: ‘clinically significant depressive illness, moderate or severe’ indicated recognition of the depression.

Measures of deprivation

The UPA scores for each electoral ward, taken from the 1991 census, were used as the index of deprivation. Patients were allocated the deprivation scores for the practice (or branch surgery) where the index consultation took place, irrespective of their own address. The non-UPA statistics from the 1991 census (Office of Population Censuses and Surveys, 1993) for car ownership and housing tenure have also been included as they appear in other measures of deprivation (Morris & Carstairs, 1991).

Data analysis

For each practice, indirect standardisation was used to adjust for age (in 10-year age bands) and gender differences in practice samples. Standardised practice prevalence ratios of probable depression were calculated by dividing the number of observed by expected cases. The relation between crude and standardised practice case prevalence and UPA score was modelled using linear regression.

At the patient level, two analyses were carried out, one using 'probable cases' examined prevalence and the other using 'possible cases' (including all probable cases) examined outcome. Multiple logistic regression was used to estimate odds ratios (ORs) for risk of probable depression. A series of models were constructed with probable depression as the dependent variable. Individual covariates of gender, age and employment status were added to the model. Odds ratios for the UPA are presented for an increase of 10 points. The final model included all possible two-way interactions with backward stepwise selection to remove non-significant terms at $P < 0.05$. Multiple logistic regression was used in the 'possible cases' to identify variables predicting 'remission' and 'improvement'. The following covariates were entered: age, gender and employment status, recognition of depression by the practitioner, depression and anxiety scores at index consultation, practice UPA score. The analyses were controlled for fund-holding and mental health resources. Interaction terms were not explored. Backward stepwise selection was used to remove non-significant variables at $P < 0.05$. Statistical procedures were performed in SPSS 6.1.2 for Windows.

RESULTS

Patients

Records of questionnaire completion were made in the second and fourth phases of screening and the average response rate was 89%, i.e. 11% of eligible patients were unable or unwilling to complete the screening instruments. There were no significant associations between practice UPA score and response rate (Spearman's correlation coefficient, 0.05, $P=0.74$ and -0.04 , $P=0.75$).

Of the patients, 12 168 (66.1%) were female and 6246 (33.9%) were male. The youngest patient was 16 and there were 194 aged between 85 and 94 years. The number of possible cases was 3668 (19.9%) patients and of these 2299 (62.7%) returned a HAD at 6 weeks and 1956 (53.3%) at 6 months. Respondents at 6 weeks did not differ significantly from non-responders in depression or anxiety scores at the index consultation (Mann-Whitney U -test, $Z=-1.81$, $P=0.07$ and $Z=-0.84$, $P=0.40$) or UPA score (Mann-Whitney U -test, $Z=-0.298$, $P=0.77$). Neither did they differ in the eventual HAD scores at 6 months' follow-up (Mann-Whitney U -test, $Z=-0.475$, $P=0.64$ and $Z=-0.841$, $P=0.40$). However, women were more likely to return questionnaires at 6 week follow-up ($\chi^2=8.62$, d.f.=1, $P=0.003$), and the response rate was positively associated with age at both assessments. At 6 months, patients from high UPA practice areas were less likely to return the questionnaire (Mann-Whitney U -test, $Z=-0.287$,

$P=0.004$), possibly confounded by greater geographical mobility.

Prevalence of depressive symptoms and deprivation scores of the practice location

The mean practice prevalence of probable depression was 7.2% (s.d. 2.6%), ranging from 2.4% to 13.7% ($\chi^2=169.8$, d.f.=57, $P < 0.0001$). The regression of crude practice prevalence rates ($n=58$ surgery addresses) against UPA score gives $\beta=0.10$ (95% CI 0.07–0.13) and $r^2=45.3\%$. The regression of standardised probable case ratios against UPA gives $\beta=1.43$ (%) (95% CI 1.03–1.82), $r^2=48.3\%$ (Fig. 1). Analysis excluding the two practices with extreme UPA scores reduced the attributable variance only slightly to $r^2=45.2\%$.

Each of the eight variables of the UPA score together with car ownership and housing tenure was examined separately for its correlation with the standardised probable case ratios (Table 1). Geographical mobility (change of address in the past year) was the only variable that did not correlate significantly. The percentage of households with two or more cars was the most highly correlated single variable ($r_s=-0.70$) with unemployment close behind (0.69).

Figure 2 shows frequency curves for practices falling above and below the median UPA score. The correlation of probable cases with deprivation appears to be due to an increase across the range of mild to moderate severity (D score 6–17) rather than to a small excess of severe cases




Fig. 1 Age- and gender-adjusted prevalence of high depression scores on the Hospital Anxiety and Depression (HAD) scale ($HAD-D \geq 11$) in 58 practice sites showing association with the Jarman under-privileged area (UPA) score of the electoral ward in which the practice is located.

or to a right shift in the whole population (Anderson *et al*, 1993).

Associations between individual characteristics and depression

All age groups from 25 years to 54 years had significantly raised ORs for depression prevalence compared to the age group 16–24 years, and the ORs fell below 1.0 (but not significantly) for all groups over 65 years. Exactly 7.1% of both females and males were cases and there was therefore no association with gender. The unemployed, temporarily away from work or permanently unable to work, and those looking after home and family, were significantly more likely to be depressed than those in employment (Table 2). The retired and student groups were at lower risk than employees.

The relationship between the two main variables was further analysed to examine the effect of practice UPA score on the odds of an individual patient being a case. Here, the UPA score of the patient's practice becomes a variable belonging to the patient. Tables 3 and 4 show the effect of UPA score in these models. The important result is that individual patient variables account for some, but not all of the variance and the UPA score remained significant after adjusting for these effects. The odds of being a case increase by 1.10 (1.06 to 1.13) for every 10 UPA points, or 1.37 (1.24 to 1.52) for 34 UPA points (2 standard deviations).

Unemployed women were less likely to be depressed than unemployed men. The self-employed, those permanently unable to work or looking after home and family were less likely to be depressed if they were older. There were no significant interaction terms with UPA score and employment status, suggesting that the effects of unemployment were as important in residents of socio-economically deprived areas as in wealthier parts.

Predictors of outcome in possible depression

The variables that significantly predicted outcome in the logistic regression models are shown in Table 5. Higher UPA scores most strongly predicted poor outcome at both 6 weeks and 6 months, for both remission and improvement. Remission was predicted by a lower initial severity of depression and anxiety and being a student.

Table 1 Spearman's correlation coefficients for the Jarman under-privileged area (UPA) score, un-transformed constituent census variables, housing tenancy and car ownership with standardised practice probable case ratios

	Correlation r_s	P value
Jarman UPA score	0.71	<0.0005
Unemployed	0.69	<0.0005
One-parent households	0.67	<0.0005
Unskilled	0.61	<0.0005
Overcrowded households	0.60	<0.0005
Children under 5 years	0.39	0.002
Ethnic minority households	0.35	0.007
Elderly living alone	0.35	0.007
Change of address in past year	0.20	0.13
Census variables		
Not owner-occupier	0.46	<0.0005
% households with no car	0.69	<0.0005
% households with two or more cars	-0.70	<0.0005




Fig. 2 Hospital Anxiety and Depression (HAD) D scores for patients attending practices in the upper and lower quartiles ranked by Jarman under-privileged area (UPA) score; ·····, least-deprived quartile; —, most-deprived quartile

Table 2 Odds ratios for risk of probable caseness by individual employment status

	Patients	Probable cases (HAD-D ≥ 11)	
	n (%)	n (%)	OR (95% CI)
Employee	6972 (37.9)	394 (5.7)	1
Self-employed	1109 (6.0)	66 (6.0)	1.06 (0.81–1.38)
Looking after home and family	2325 (12.6)	212 (9.1)	1.68 (1.41–1.99)
Retired	4588 (24.9)	208 (4.5)	0.79 (0.67–0.94)
Unemployed	764 (4.1)	116 (15.2)	2.99 (2.39–3.73)
Student	1036 (5.6)	33 (3.2)	0.55 (0.38–0.79)
Permanently unable to work	569 (3.1)	132 (23.2)	5.04 (4.05–6.28)
Temporarily away from work	555 (3.0)	104 (18.7)	3.84 (3.03–4.86)
Missing data	495 (2.7)	42 (8.5)	1.55 (1.11–2.16)

HAD–D, Hospital Anxiety and Depression scale, depression score.

Table 3 The effect of the Jarman under-privileged area (UPA) score on the odds of being a case, adjusted for the intervention of the randomised controlled trial from which the sample was derived, the phase at which the patient was screened, gender, age, and employment status ($n=18\ 414$)

Model	Variable	UPA score (10 points)	
		OR (95% CI)	P value
M1	UPA score alone	1.13 (1.09–1.16)	<0.00005
M2	M1+intervention group, phase and group by phase interaction	1.12 (1.09–1.16)	<0.00005
M3	M2+gender	1.12 (1.09–1.16)	<0.00005
M4	M3+age	1.12 (1.09–1.15)	<0.00005
M5	M4+employment status	1.09 (1.06–1.13)	<0.00005
M6	Final model (see Table 4)	1.10 (1.07–1.13)	<0.00005

Table 4 Adjusted odds ratios (95% CI) for each significant variable in the final model

Variable	Odds ratio	(95% CI)	P value
UPA score (10 points)	1.10	(1.06–1.13)	<0.00005
Age (years)	1.01	(1.00–1.01)	0.0465
Employee	1		
Self-employed	2.77	(1.06–7.19)	0.0368
Interaction with age	0.98	(0.96–1.00)	0.0456
Looking after home and family	2.80	(1.66–4.73)	0.0001
Interaction with age	0.99	(0.97–1.00)	0.0351
Retired	7.59	(2.53–22.8)	0.0003
Interaction with age	0.97	(0.95–0.98)	<0.00005
Unemployed	3.73	(2.83–4.91)	<0.00005
Interaction with gender (female)	0.54	(0.36–0.82)	0.0034
Student	0.57	(0.39–0.84)	0.0040
Permanently unable to work	16.6	(6.62–41.8)	<0.00005
Interaction with age	0.97	(0.96–0.99)	0.0050
Temporarily away from work	3.71	(2.93–4.71)	<0.00005

UPA, Jarman under-privileged area.

Improvement was independent of initial severity of depression but adversely influenced by anxiety.

Recognition of depression by the GP predicted remission (but not improvement) at 6 weeks but not at 6 months. No differential effect was found for gender. Older patients were less likely to have improved at 6 weeks and the retired group at 6 months. Thus the most consistent predictors of outcome were the socio-economic variables of area deprivation and individual employment status.

DISCUSSION

These results support an influence of socio-economic deprivation on both the

prevalence and persistence of depression among general practice attenders that is of practical consequence. The UPA score accounted for almost half the inter-practice variation in prevalence – an effect produced by an increase in mild to moderate cases (D score 6–17). Individual characteristics of unemployment and car ownership were of comparable strength but the UPA score remained the strongest predictor of depression and of outcome up to 6 months later.

The general practices were representative of Hampshire, being larger than those in England as a whole and generally more affluent than inner-city or northern areas. However, they were situated in areas with a wide range of deprivation scores. The study was carried out within 4–5 years of

the 1991 census from which UPA scores were derived. The definition of depression was not dependent on the accuracy of general practitioners' diagnoses, thus avoiding potential bias.

Limitations of the study

It was logistically impossible to carry out a diagnostic interview of more than 18 000 patients. We have therefore studied the prevalence of self-reported depressive symptoms rather than of an operationally defined diagnosis. Nevertheless, in primary care studies screening for depressive symptoms rather than syndromes is an appropriate strategy, since sub-syndromal major depression has been shown to contribute significantly to the public health burden (Hays *et al*, 1995; Olfson *et al*, 1996). For this purpose, the choice of the HAD appears to have been appropriate despite its modest sensitivity and specificity for DSM-III (American Psychiatric Association, 1980) mood disorders.

The sample of practices was broadly representative but it was not randomly selected since it depended upon at least one practitioner in each practice agreeing to take part. No inferences can be made from this study about depressive symptoms in the community at large nor about patients seen at home. However, it is known that home visits account for only 10% of doctor-patient contacts (McCormick *et al*, 1995) apart from elderly patients, where they are more frequent (35% at 75–84 years and 65% over 85 years), so the study results may not hold true for the elderly infirm.

The rate of follow-up was lower than we had expected and was affected by age, gender and UPA score, but not HAD scores, on at least one time point. It seems unlikely that the association we found between HAD and UPA scores was critically influenced by non-response in the absence of an association between response rate and HAD score. However, we cannot rule out the possibility that this association was affected by non-responders.

The measure of area deprivation was not exact because some patients might have lived in an adjacent electoral ward, not in that of the practice. This would have introduced a conservative bias into the associations. The UPA score has been criticised on a number of counts (Morris & Carstairs, 1991) and some have suggested that complex indices are unnecessary because

Table 5 Factors at baseline that significantly predicted 6-week and 6-month outcome in those with possible depression

	Outcome measure			
	Remission		Improvement	
	OR (95% CI)	P value	OR (95% CI)	P value
<i>6 weeks' follow-up</i>				
UPA score (10 points)	0.95 (0.90–0.99)	0.0311	0.90 (0.84–0.97)	0.0069
HAD, depression score	0.75 (0.71–0.79)	<0.0001	–	
HAD, anxiety score	0.96 (0.94–0.99)	0.0032	0.97 (0.94–1.00)	0.0469
Age (years)	–		0.99 (0.98–1.00)	0.0446
Rated as depressed by GP	1.31 (1.06–1.61)	0.0126	–	
Looking after home and family	–		0.65 (0.44–0.95)	0.0274
Permanently unable to work	0.56 (0.36–0.87)	0.0101	0.44 (0.22–0.88)	0.0202
Temporarily away from work	–		0.49 (0.24–0.98)	0.0434
<i>6 months' follow-up</i>				
UPA score (10 points)	0.92 (0.87–0.97)	0.0021	0.92 (0.86–0.98)	0.0138
HAD, depression score	0.82 (0.78–0.85)	<0.0001	–	
Looking after home and family	0.74 (0.56–0.97)	0.0318	0.63 (0.44–0.89)	0.0084
Retired	–		0.51 (0.37–0.70)	0.0000
Unemployed	0.66 (0.43–1.01)	0.0535	0.49 (0.28–0.86)	0.0137
Student	2.44 (1.40–4.27)	0.0017	–	
Permanently unable to work	0.41 (0.26–0.65)	0.0002	0.29 (0.15–0.55)	0.0002

UPA, Jarman under-privileged area; HAD, Hospital Anxiety and Depression scale; GP, general practitioner.

unemployment rates alone are an adequate measure of deprivation (Payne *et al*, 1993). We found positive correlations between depression and most component variables of the index but the total score showed a higher correlation than any single variable, with only unemployment and car ownership approaching it. The precise choice of index is probably irrelevant since they are all highly correlated (Morris & Carstairs, 1991).

We did not measure causes of poverty other than unemployment at the individual level so we have insufficient data fully to separate out the effects of area and individual deprivation, but correcting for individual unemployment only slightly reduced the effect of UPA score. It seems, therefore, that there may be an effect of living in a deprived area which may have an influence over and above that of individual deprivation.

Interpretation of the findings

Given the strength of these correlations it would not be unreasonable to assume that they are a reflection of those in the general population, even though they were measured in primary care attenders. If so,

the associations we have demonstrated may come about through two main routes (Stansfield *et al*, 1998) and our results do not allow us to distinguish between them. First (the social cause hypothesis), living in a deprived area may cause depression perhaps with material insecurity, loss of social cohesion or a non-economic predisposing factor, such as certain types of early life experience acting as intervening variables (Power & Hertzman, 1997). That the environment may play a part is suggested by studies that have reported reductions in mental illness after improvement in the urban environment (Halpern, 1995). Second, there may be social selection with depressed individuals moving into deprived areas – as occurs with some suffering from schizophrenia (Sloggett & Joshi, 1994). The relative influence of social cause and social selection may vary with the economic climate (Hammarstrom & Janlert, 1997).

This is the first time that area deprivation has been examined as a predictor of the persistence of depression in consulting patients. Support for these findings can, however, be found in studies of individual risk factors where chronic social difficulties, low household income and perceived

financial strain have been associated with the maintenance of depression (Brown & Moran, 1997; Ronalds *et al*, 1997; Weich *et al*, 1997).

The difference in the distribution of D scores in practices in the top and bottom quartile of the deprivation score is the kind described by Anderson *et al* (1993) as showing the presence of “a subset of individuals [who are] particularly exposed or susceptible [to a postulated cause acting at an individual level]”. This distinguishes it from a cause acting on a population, all of whom are equally susceptible, which would produce a right shift in the entire distribution.

Implications of the results

In practical terms, our findings have two implications for health policy. First, the emphasis on enhancing provision to deprived areas is correct, as anticipated, but there is no threshold above which the effects of deprivation begin to become apparent. Thus, any model of deprivation payments that has a built-in threshold will disadvantage those areas just below it. Since deprivation accounts for such a large proportion of the depression workload this may have a measurable effect on the equity of primary mental health care provision. Second, strategies to reduce the public health burden of depression might usefully combine population-based and high-risk approaches: for example, strong national environmental and employment policies in combination with health care that acknowledges the adverse influence of living in a deprived area. While individuals may benefit from specific treatments for depression there is little evidence that even their most effective use, without other measures, could significantly reduce the public health burden of the condition.

ACKNOWLEDGEMENTS

The study was funded by grants from the Medical Research Council, the South and West Regional Research and Development Committee and the Southampton Community Health Service NHS Trust. We thank Professor Brian Jarman for providing the scores for Hampshire and Professor T. Kendrick for helpful comments on the manuscript.

REFERENCES

American Psychiatric Association (1980) *Diagnostic and Statistical Manual of Mental Disorders* (3rd edn) (DSM-III). Washington, DC:APA.

Acheson, O. (1998) *Independent Inquiry into Inequalities in Health*. London: HMSO.

Anderson, J., Huppert, F. & Rose, G. (1993) Normality, deviance and minor psychiatric morbidity in the community. A population-based approach to General Health Questionnaire data in the Health and Lifestyle Survey. *Psychological Medicine*, **23**, 475–485.

Brown, G. W. & Moran, P. M. (1997) Single mothers, poverty and depression. *Psychological Medicine*, **27**, 21–33.

Eachus, J., Williams, M., Chan, P., et al (1996) Deprivation and cause specific morbidity: evidence from the Somerset and Avon survey of health. *British Medical Journal*, **312**, 287–292.

Halpern, D. (1995) *Mental Health and Built Environment. More than Bricks and Mortar*. London: Taylor & Francis.

Hammarstrom, A. & Janlert, U. (1997) Nervous and depressive symptoms in a longitudinal study of youth unemployment – selection or exposure? *Journal of Adolescence*, **20**, 293–305.

Hays, R. D., Wells, K. B., Sherbourne, C. D., et al (1995) Functioning and well-being outcomes of patients with depression compared with chronic general medical illnesses. *Archives of General Psychiatry*, **52**, 11–19.

Herrman, C. (1997) International experiences with the hospital anxiety and depression scale – a review of validation data and clinical results. *Journal of Psychosomatic Research*, **42**, 17–41.

Jarman, B. (1983) Identification of under-privileged areas. *British Medical Journal*, **286**, 1705–1709.

McCormick, A., Charlton, J. & Fleming, D. (1995) Who sees their general practitioner and for what reason? *Health Trends*, **27**, 34–35.

Meltzer, H., Gill, B., Petticrew, M., et al (1995) *The Prevalence of Psychiatric Morbidity among Adults Living in Private Households*. OPCS Surveys of Psychiatric Morbidity in Great Britain. Report 1. London: HMSO.

Morris, R. & Carstairs, V. (1991) Which deprivation? A comparison of selected deprivation indexes. *Journal of Public Health Medicine*, **13**, 318–326.

Office of Population Censuses and Surveys (1993) *1991 Census. County Report: Hampshire (Part 1) Volume 1*. CEN 91 CR 19. London: HMSO.

Olfson, M., Broadhead, W. E., Weissman, M. M., et al (1996) Subthreshold psychiatric symptoms in a primary care group practice. *Archives of General Psychiatry*, **53**, 880–886.

Payne, J. N., Coy, J., Milner, P. C., et al (1993) Are deprivation indicators a proxy for morbidity? A comparison of the prevalence of arthritis, depression, dyspepsia, obesity and respiratory symptoms with unemployment rates and Jarman scores. *Journal of Public Health Medicine*, **15**, 161–171.

Power, C. & Hertzman, C. (1997) Social and biological pathways linking early life and adult disease. *British Medical Bulletin*, **53**, 210–221.

Ronalds, C., Creed, F., Stone, K., et al (1997) Outcome of anxiety and depressive disorders in primary care. *British Journal of Psychiatry*, **171**, 427–433.

CLINICAL IMPLICATIONS

- In primary care attenders, socio-economic deprivation accounted for half the large inter-practice variation in the prevalence of depression and predicted the persistence of depression over 6 months.
- The effect of deprivation appears to be exerted through its influence on vulnerable individuals rather than the whole population.
- Efforts at reducing the public health burden of depression should include social strategies as well as medical care.

LIMITATIONS

- The practices taking part were representative of Hampshire practices in all measured respects, but were not randomly selected.
- Cases were defined by scores on the Hospital Anxiety and Depression scale, not operational diagnostic criteria.
- The results from 6-week and 6-month follow-ups were based on modest response rates.

K. OSTLER, MRCPsych, C. THOMPSON, FRCP, A.-L. K. KINMONTH, FRCGP, R. C. PEVELER, FRCPsych, L. STEVENS, MRCPsych, A. STEVENS, FRCP, Community Based Clinical Sciences Research Division, Faculty of Medicine, Health and Biological Sciences, University of Southampton

Correspondence: C. Thompson, Department of Psychiatry, Royal South Hants Hospital, Brinton's Terrace, Southampton SO14 0YG

(First received 19 November 1999, final revision 5 April 2000, accepted 19 June 2000)

Shepherd, M., Cooper, B., Brown, A. C., et al (1966) *Psychiatric Illness in General Practice*. London: Oxford University Press.

Sloggett, A. & Joshi, H. (1994) Higher mortality in deprived areas: community or personal disadvantage? *British Medical Journal*, **309**, 1470–1474.

Stansfeld, S. A. & Marmot, M. G. (1992) Social class and minor psychiatric disorder in British civil servants: a validated screening survey using the General Health Questionnaire. *Psychological Medicine*, **22**, 739–749.

—, **Head, J. & Marmot, M. G. (1998)** Explaining social class differences in depression and well being. *Social Psychiatry and Psychiatric Epidemiology*, **33**, 1–9.

Thompson, C., Kinmonth, A.-L., Stevens, L. et al (2000) Effects of a clinical-practice guideline and practice-based education on detection and outcome of depression in primary care: Hampshire Depression Project, randomised controlled trial. *Lancet*, **355**, 185–191.

Thornicroft, G., Margolius, O. & Jones, D. (1992) The TAPS Project. 6: New long-stay psychiatric patients and social deprivation. *British Journal of Psychiatry*, **161**, 621–624.

Upadhyaya, A. K. & Stanley, I. (1993) Hospital Anxiety and Depression Scale. *British Journal of General Practice*, **43**, 349–350.

Weich, S., Churchill, R., Lewis, G., et al (1997) Do socio-economic risk factors predict the incidence and maintenance of psychiatric disorder in primary care? *Psychological Medicine*, **27**, 73–80.

Wilkinson, M. J. B. & Barczak, P. (1988) Psychiatric screening in general practice: comparison of the General Health Questionnaire and the Hospital Anxiety Depression scale. *Journal of the Royal College of General Practitioners*, **38**, 311–313.

Zigmond, A. S. & Snaith, R. P. (1983) The Hospital Anxiety and Depression Scale. *Acta Psychiatrica Scandinavica*, **67**, 361–370.